


assachusetts Division of Fisheries & Wildlife Route 135, Westborough, MA 01581 tel: (508) 389-6360; fax: (508) 389-7891 www.nhesp.org

Description: The Orange Sallow is a noctuid moth with orange forewings spotted with black, the wing margins pink in the male; the hind wings are tan, the wing margins shaded with pink in both sexes. Wingspan is 24-28 mm.

Habitat: The Orange Sallow Moth inhabits xeric and open oak woodland on rocky uplands (ridges, hilltops, and steep slopes), and the edges of old fields, powerline cuts, and other openings within such habitat.

Life History: Adult moths fly in August. Eggs hatch soon after they are laid, and larvae feed on the flowers, seeds, and foliage of false foxgloves (*Aureolaria*) through early October. Pupae overwinter and diapause through the following spring and summer.

Range: The Orange Sallow Moth is spottily distributed from southern New England south to Florida, west to Wisconsin and Missouri. In Massachusetts this species occurs in scattered colonies from Boston west to the Berkshires, but is absent from the southeast coastal plain.



Orange Sallow Moth

Rhodoecia aurantiago

State Status: **Threatened** Federal Status: None



Photo by M.W. Nelson

Adult Flight Period in Massachusetts

		_		8																			
Jan		Feb		Mar		Apr		May		Jun		Jul		Aug		Sep		Oct		Nov		Dec	

Threats

- Habitat loss
- Fire suppression
- Invasion by exotic plants
- Introduced generalist parasitoids
- Insecticide spraying
- Clearcut timber harvest
- Excessive deer browsing of larval host plants
- Off-road vehicles
- Light pollution